

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number
WO 02/02519 A2

- (51) International Patent Classification⁷: **C07D**
- (21) International Application Number: PCT/US01/20962
- (22) International Filing Date: 29 June 2001 (29.06.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/214,831 29 June 2000 (29.06.2000) US
- (71) Applicant: **DUPONT PHARMACEUTICALS COMPANY** [US/US]; Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 (US).
- (72) Inventors: **PINTO, Donald, J., P.**; 425 Brentford Road, Kennett Square, PA 19348 (US). **QUAN, Mimi, L.**; 113 Venus Drive, Newark, DE 19711 (US).
- (74) Agent: **ROPER, David, J.**; Dupont Pharmaceuticals Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).

- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/02519 A2

(54) Title: **THROMBIN OR FACTOR Xa INHIBITORS**

(57) Abstract: This invention relates generally to heteroaryl-phenyl substituted compounds that are inhibitors of trypsin-like serine protease enzymes, especially factor Xa or thrombin, pharmaceutical compositions containing the same, and methods of using the same as anticoagulant agents for treatment and prevention of thromboembolic disorders.

TITLE

Thrombin or Factor Xa Inhibitors

5

FIELD OF THE INVENTION

This invention relates generally to heteroaryl-phenyl substituted compounds that are inhibitors of trypsin-like serine protease enzymes, especially factor Xa or thrombin, pharmaceutical compositions containing the same, and methods
10 of using the same as anticoagulant agents for treatment and prevention of thromboembolic disorders.

BACKGROUND OF THE INVENTION

Activated factor Xa, whose major practical role is the
15 generation of thrombin by the limited proteolysis of prothrombin, holds a central position that links the intrinsic and extrinsic activation mechanisms in the final common pathway of blood coagulation. The generation of thrombin, the final serine protease in the pathway to
20 generate a fibrin clot, from its precursor is amplified by formation of prothrombinase complex (factor Xa, factor V, Ca^{2+} and phospholipid). Since it is calculated that one molecule of factor Xa can generate 138 molecules of thrombin, inhibition of factor Xa may be more efficient than
25 inactivation of thrombin in interrupting the blood coagulation system.

Therefore, efficacious and specific inhibitors of factor Xa, thrombin, or both are needed as potentially valuable therapeutic agents for the treatment of
30 thromboembolic disorders. It is thus desirable to discover new factor Xa, thrombin, or both inhibitors.

SUMMARY OF THE INVENTION

Accordingly, one object of the present invention is to provide novel heteroaryl-phenyl substituted compounds that are useful as factor Xa inhibitors or pharmaceutically acceptable salts or prodrugs thereof.

It is another object of the present invention to provide pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.


It is another object of the present invention to provide a method for treating thromboembolic disorders comprising administering to a host in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.


It is another object of the present invention to provide novel compounds for use in therapy.

It is another object of the present invention to provide the use of novel compounds for the manufacture of a medicament for the treatment of thrombosis or a disease mediated by factor Xa.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[1] Thus, in an embodiment, the present invention provides a novel compound selected from the group:







or a stereoisomer or pharmaceutically acceptable salt thereof, wherein;

5


G is selected from formulas Ia-Ic:



Ia



Ib



Ic

10

ring D₁ is selected from pyridine, pyrazine, pyridazine, and pyrimidine and is substituted with 1 D_{1a} and 0-1 D_{1b};

ring D₂ is a 5-membered heteroaromatic ring system

15

comprising E, carbon atoms, and 0-3 N atoms, wherein E is selected from O, S, and N-D_{1c} and ring D₂ is substituted with 1 D_{1a} and 0-1 D_{1b};

ring D₃ is a 5-membered heteroaromatic ring system
 comprising carbon atoms and from 0-3 additional N atoms
 and ring D₃ is substituted with 1 D_{1a} and 0-1 D_{1b};

5 G¹ is selected from H, C₁₋₄ alkyl, F, Cl, Br, I, OH, OCH₃,
 OCH₂CH₃, OCH(CH₃)₂, OCH₂CH₂CH₃, CN, C(=NR⁸)NR⁷R⁹,
 NHC(=NR⁸)NR⁷R⁹, NR⁸CH(=NR⁷), NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃
 alkyl)₂, C(=NH)NH₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl), CH₂N(C₁₋₃
 alkyl)₂, CH₂CH₂NH₂, CH₂CH₂NH(C₁₋₃ alkyl), CH₂CH₂N(C₁₋₃
 10 alkyl)₂, (CR⁸R⁹)_tNR⁷R⁸, (CR⁸R⁹)_tC(O)NR⁷R⁸, and OCF₃;

D_{1a} is selected from H, C₁₋₄ alkyl, F, Cl, Br, I, OH, OCH₃,
 OCH₂CH₃, OCH(CH₃)₂, OCH₂CH₂CH₃, CN, C(=NR⁸)NR⁷R⁹,
 NHC(=NR⁸)NR⁷R⁹, NR⁸CH(=NR⁷), NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃
 15 alkyl)₂, C(=NH)NH₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl), CH₂N(C₁₋₃
 alkyl)₂, CH₂CH₂NH₂, CH₂CH₂NH(C₁₋₃ alkyl), CH₂CH₂N(C₁₋₃
 alkyl)₂, (CR⁸R⁹)_tNR⁷R⁸, (CR⁸R⁹)_tC(O)NR⁷R⁸, and OCF₃;

D_{1b} is selected from H, C₁₋₄ alkyl, F, Cl, Br, I, OH, OCH₃,
 20 OCH₂CH₃, OCH(CH₃)₂, OCH₂CH₂CH₃, CN, C(=NR⁸)NR⁷R⁹,
 NHC(=NR⁸)NR⁷R⁹, NR⁸CH(=NR⁷), NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃
 alkyl)₂, C(=NH)NH₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl), CH₂N(C₁₋₃
 alkyl)₂, CH₂CH₂NH₂, CH₂CH₂NH(C₁₋₃ alkyl), CH₂CH₂N(C₁₋₃
 alkyl)₂, (CR⁸R⁹)_tNR⁷R⁸, (CR⁸R⁹)_tC(O)NR⁷R⁸, and OCF₃;

25 D_{1c} is selected from H, C₁₋₄ alkyl, OCH₃, OCH₂CH₃, OCH(CH₃)₂,
 OCH₂CH₂CH₃, NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, C(=NH)NH₂,
 CH₂NH₂, CH₂NH(C₁₋₃ alkyl), CH₂N(C₁₋₃ alkyl)₂, CH₂CH₂NH₂,
 CH₂CH₂NH(C₁₋₃ alkyl), CH₂CH₂N(C₁₋₃ alkyl)₂, (CR⁸R⁹)_tNR⁷R⁸,
 30 (CR⁸R⁹)_tC(O)NR⁷R⁸, and OCF₃;

G^2 is absent or is selected from CH_2 , $C(O)$, O , NR^3 , $S(O)_p$,
 CH_2CH_2 , $C(O)CH_2$, $CH_2C(O)$, OCH_2 , CH_2O , NR^3CH_2 , CH_2NR^3 ,
 $S(O)_pCH_2$, $CH_2S(O)_p$, $CH_2CH_2CH_2$, $C(O)CH_2CH_2$, $CH_2C(O)CH_2$,
5 $CH_2CH_2C(O)$, OCH_2CH_2 , CH_2OCH_2 , CH_2CH_2O , $NR^3CH_2CH_2$, $CH_2NR^3CH_2$,
 $CH_2CH_2NR^3$, $S(O)_pCH_2CH_2$, $CH_2S(O)_pCH_2$, and $CH_2CH_2S(O)_p$;

G^3 is phenyl, naphthyl, or a 5-10 membered heteroaryl
 consisting of carbon atoms and from 1-3 heteroatoms
 10 selected from N, O, and S;

L_n is a linker which is absent or is selected from O, S,
 $S(O)_2$, CH_2 , $*NHC(O)$, $*C(O)NH$, $*S(O)_2NH$, $*NHS(O)_2$,
 $*CH_2NHC(O)$, $*CH(R^a)NHC(O)$, $*CH_2NHC(O)CH_2$, and
 15 $*CH(R^a)NHC(O)CH_2$, provided that L_n and M do not form an
 O-N or S-N bond and the * indicates where L_n is bonded
 to G;

M^1 is absent or is selected from CHR , O , and NR^2 ;
 20

M^2 is N or CR^f ;

M^3 is N or CR^d ;

25 provided that only one of M^2 and M^3 is N;

M^4 is selected from NR^2 , CR^f , and $C(O)$;

R^a is selected from $C(O)C(O)OR^3$, $C(O)C(O)NR^2R^{2a}$, and $C(O)-A$;
 30

R^b is selected from H, R, phenyl, C_{1-10} alkyl, and C_{2-5} alkenyl;

R^c is selected from H and C_{1-6} alkyl;

5

alternatively, R^b and R^c together are $-(CH_2)_4-$;

R^d is selected from H, F, and Cl;

10 R^e is selected from H, $N(CH_3)(CH_2CO_2H)$ and S-(5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^4);

15 alternatively, R^d and R^e combine to form $-NR^3-C(O)-C(R^1gR^3)-NR^3-$ or $-N=CR^2-NR^3-$;

R^f is selected from H, F, and Cl;

20 alternatively, R^e and R^f combine to form $-NR^3-C(R^1gR^3)-C(O)-NR^3-$ or $-NR^3-CR^2=N-$;

R^g is selected from H, CH_2OR^3 , $CH_2C(O)OR^3$, C_{1-4} alkyl, $C(O)NH_2$, and NH_2 ;

25

R^h is selected from H, CH_2 -phenyl, CH_2CH_2 -phenyl, and $CH=CH$ -phenyl;

R^i is selected from $SO_2CH_2C(O)OR^3$, $C(O)CH_2C(O)OR^3$, and $C(O)OR^3$;

30

R is selected from H, Cl, F, Br, I, $(\text{CH}_2)_t\text{OR}^3$, C_{1-4} alkyl, benzyl, OCF_3 , CF_3 , $\text{C}(\text{O})\text{NR}^7\text{R}^8$, $(\text{CH}_2)_t\text{NR}^2\text{SO}_2\text{-C}_{1-4}$ alkyl, and $(\text{CR}^8\text{R}^9)_t\text{NR}^7\text{R}^8$;

- 5 Z is selected from a $(\text{CR}^8\text{R}^9)_{1-4}$, $(\text{CR}^8\text{R}^9)_r\text{O}(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{NR}^3(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{NR}^3\text{C}(=\text{CHR}^8)(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{C}(\text{O})(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{C}(\text{O})\text{O}(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{OC}(\text{O})(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{C}(\text{O})\text{NR}^3(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{NR}^3\text{C}(\text{O})(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{OC}(\text{O})\text{O}(\text{CR}^8\text{R}^9)_r$,
 10 $(\text{CH}_2)_r\text{OC}(\text{O})\text{NR}^3(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{NR}^3\text{C}(\text{O})\text{O}(\text{CR}^8\text{R}^9)_r$, $(\text{CH}_2)_r\text{NR}^3\text{C}(\text{O})\text{NR}^3(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{S}(\text{O})_p(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{S}(\text{O})_2(\text{CH}=\text{CH})$, $(\text{CCR}^8\text{R}^9)_r\text{SO}_2\text{NR}^3(\text{CR}^8\text{R}^9)_r$, $(\text{CR}^8\text{R}^9)_r\text{NR}^3\text{SO}_2(\text{CR}^8\text{R}^9)_r$, and $(\text{CR}^8\text{R}^9)_r\text{NR}^3\text{SO}_2\text{NR}^3(\text{CR}^8\text{R}^9)_r$,
 provided that Z does not form a N-N, N-O, N-S, NCH_2N ,
 15 NCH_2O , or NCH_2S bond with the groups to which Z is attached;

- R^{1a} is selected from H, $-(\text{CH}_2)_r\text{-R}^{1b}$, $-\text{CH}=\text{CH-R}^{1b}$, $\text{NCH}_2\text{R}^{1c}$, $\text{OCH}_2\text{R}^{1c}$, $\text{SCH}_2\text{R}^{1c}$, $\text{NH}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1b}$, $\text{O}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1b}$,
 20 $\text{S}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1b}$, $\text{S}(\text{O})_p(\text{CH}_2)_r\text{R}^{1d}$, $\text{O}(\text{CH}_2)_r\text{R}^{1d}$, $\text{NR}^3(\text{CH}_2)_r\text{R}^{1d}$, $\text{OC}(\text{O})\text{NR}^3(\text{CH}_2)_r\text{R}^{1d}$, $\text{NR}^3\text{C}(\text{O})\text{NR}^3(\text{CH}_2)_r\text{R}^{1d}$, $\text{NR}^3\text{C}(\text{O})\text{O}(\text{CH}_2)_r\text{R}^{1d}$, and $\text{NR}^3\text{C}(\text{O})(\text{CH}_2)_r\text{R}^{1d}$, provided that R^{1a} forms other than an N-halo, N-N, N-S, N-O, or N-CN bond;

- 25 R^{1b} is selected from H, C_{1-3} alkyl, F, Cl, Br, I, $-\text{CN}$, $-\text{CHO}$, $(\text{CF}_2)_r\text{CF}_3$, $(\text{CH}_2)_r\text{OR}^2$, NR^2R^{2a} , $\text{C}(\text{O})\text{R}^{2c}$, $\text{OC}(\text{O})\text{R}^2$, $(\text{CF}_2)_r\text{CO}_2\text{R}^{2a}$, $\text{S}(\text{O})_p\text{R}^{2b}$, $\text{NR}^2(\text{CH}_2)_r\text{OR}^2$, $\text{C}(=\text{NR}^{2c})\text{NR}^2\text{R}^{2a}$, $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$, $\text{NR}^2\text{C}(\text{O})\text{NHR}^{2b}$, $\text{NR}^2\text{C}(\text{O})_2\text{R}^{2a}$, $\text{OC}(\text{O})\text{NR}^{2a}\text{R}^{2b}$, $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$, $\text{C}(\text{O})\text{NR}^2(\text{CH}_2)_r\text{OR}^2$, $\text{SO}_2\text{NR}^2\text{R}^{2a}$, $\text{NR}^2\text{SO}_2\text{R}^{2b}$, C_{3-6}

carbocycle substituted with 0-2 R^{4a} , and 5-10 membered heterocycle consisting of carbon atoms and from 1-4 heteroatoms selected from the group consisting of N, O, and S(O), substituted with 0-2 R^{4a} , provided that R^{1b} forms other than an N-halo, N-N, N-S, N-O, or N-CN bond;

R^{1c} is selected from H, $CH(CH_2OR^2)_2$, $C(O)R^{2c}$, $C(O)NR^2R^{2a}$, $S(O)R^{2b}$, $S(O)_2R^{2b}$, and $SO_2NR^2R^{2a}$;

10

R^{1d} is selected from C_{3-13} carbocycle substituted with 0-2 R^{4a} , and 5-13 membered heterocycle consisting of carbon atoms and from 1-4 heteroatoms selected from the group consisting of N, O, and S(O), substituted with 0-2 R^{4a} , provided that R^{1d} forms other than an N-N, N-S, or N-O bond;

15

R^{1g} is selected from H, C_{1-6} alkyl, and C_{1-6} alkyl substituted with A;

20

R^2 , at each occurrence, is selected from H, CF_3 , C_{1-6} alkyl, benzyl, C_{3-6} carbocyclic residue substituted with 0-2 R^{4b} , and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b} ;

25

R^{2a} , at each occurrence, is selected from H, CF_3 , C_{1-6} alkyl, benzyl, C_{3-6} cycloalkylmethyl substituted with 0-2 R^{4b} , C_{3-6} carbocyclic residue substituted with 0-2 R^{4b} , and 5-6 membered heterocyclic system containing from 1-4

30

heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b};

R^{2b}, at each occurrence, is selected from CF₃, C₁₋₄ alkoxy, C₁₋₆ alkyl, benzyl, C₃₋₆ carbocyclic residue substituted with 0-2 R^{4b}, and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b};

10 R^{2c}, at each occurrence, is selected from CF₃, OH, C₁₋₄ alkoxy, C₁₋₆ alkyl, benzyl, C₃₋₆ carbocyclic residue substituted with 0-2 R^{4b}, and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with
15 0-2 R^{4b};

alternatively, R² and R^{2a}, together with the atom to which they are attached, combine to form a 5 or 6 membered saturated, partially saturated or unsaturated ring
20 substituted with 0-2 R^{4b} and containing from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

R³, at each occurrence, is selected from H, C₁₋₄ alkyl, and
25 phenyl;

R^{3a}, at each occurrence, is selected from H, C₁₋₄ alkyl, and phenyl;

30 R^{3b}, at each occurrence, is selected from H, C₁₋₄ alkyl, and phenyl;

R^{3c} , at each occurrence, is selected from C_{1-4} alkyl, and phenyl;

5 R^{3d} , at each occurrence, is selected from H, C_{1-4} alkyl, C_{1-4} alkyl-phenyl, and $C(=O)R^{3c}$;

A is selected from:

C_{3-10} carbocyclic residue substituted with 0-2 R^4 , and
 10 5-12 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^4 ;

A^1 is H or A;

15

alternatively, A and A^1 and the carbon to which they are attached combine to form fluorene;

A^2 is selected from H, A, and $CHAA^4$;

20

A^3 is selected from H, A, C_{1-4} alkyl, and $-(CH_2)_xNR^2R^{2a}$;

A^4 is H or A;

25 B is selected from: H, Y, and X-Y, provided that Z and B are attached to different atoms on A;

X is selected from $-(CR^2R^{2a})_{1-4}-$, $-CR^2(CR^2R^{2b})(CH_2)_t-$, $-C(O)-$,
 $-C(=NR^{1c})-$, $-CR^2(NR^{1c}R^2)-$, $-CR^2(OR^2)-$, $-CR^2(SR^2)-$,
 30 $-C(O)CR^2R^{2a}-$, $-CR^2R^{2a}C(O)-$, $-S-$, $-S(O)-$, $-S(O)_2-$,
 $-SCR^2R^{2a}-$, $-S(O)CR^2R^{2a}-$, $-S(O)_2CR^2R^{2a}-$, $-CR^2R^{2a}S-$,

$-\text{CR}^2\text{R}^{2a}\text{S}(\text{O})-$, $-\text{CR}^2\text{R}^{2a}\text{S}(\text{O})_2-$, $-\text{S}(\text{O})_2\text{NR}^2-$, $-\text{NR}^2\text{S}(\text{O})_2-$,
 $-\text{NR}^2\text{S}(\text{O})_2\text{CR}^2\text{R}^{2a}-$, $-\text{CR}^2\text{R}^{2a}\text{S}(\text{O})_2\text{NR}^2-$, $-\text{NR}^2\text{S}(\text{O})_2\text{NR}^2-$,
 $-\text{C}(\text{O})\text{NR}^2-$, $-\text{NR}^2\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^2\text{CR}^2\text{R}^{2a}-$, $-\text{NR}^2\text{C}(\text{O})\text{CR}^2\text{R}^{2a}-$,
 $-\text{CR}^2\text{R}^{2a}\text{C}(\text{O})\text{NR}^2-$, $-\text{CR}^2\text{R}^{2a}\text{NR}^2\text{C}(\text{O})-$, $-\text{NR}^2\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})\text{NR}^2-$,
5 $-\text{NR}^2\text{C}(\text{O})\text{NR}^2-$, $-\text{NR}^2-$, $-\text{NR}^2\text{CR}^2\text{R}^{2a}-$, $-\text{CR}^2\text{R}^{2a}\text{NR}^2-$, O ,
 $-\text{CR}^2\text{R}^{2a}\text{O}-$, and $-\text{OCR}^2\text{R}^{2a}-$;

Y is selected from:

C₃₋₁₀ carbocyclic residue substituted with 0-2 R^{4a}, and
 10 5-12 membered heterocyclic system containing from 1-4
 heteroatoms selected from the group consisting of N, O, and
 S substituted with 0-2 R^{4a};

alternatively, Z-A-B combine to form S-C₁₋₆ alkyl;

15

R⁴, at each occurrence, is selected from H, =O, (CH₂)_rOR²,
 (CH₂)_rF, (CH₂)_rCl, (CH₂)_rBr, (CH₂)_rI, C₁₋₄ alkyl,
 (CH₂)_rCN, (CH₂)_rNO₂, (CH₂)_rNR²R^{2a}, C(O)R^{2c}, NR²C(O)R^{2b},
 C(O)NR²R^{2a}, NR²C(O)NR²R^{2a}, C(=NR²)NR²R^{2a},
 20 C(=NS(O)₂R⁵)NR²R^{2a}, NHC(=NR²)NR²R^{2a}, C(O)NHC(=NR²)NR²R^{2a},
 SO₂NR²R^{2a}, NR²SO₂NR²R^{2a}, NR²SO₂-C₁₋₄ alkyl, NR²SO₂R⁵,
 S(O)_pR⁵, (CF₂)_rCF₃, (CH₂)_r-CF₃, NCH₂R^{1c}, OCH₂R^{1c}, SCH₂R^{1c},
 N(CH₂)₂(CH₂)_tR^{1b}, O(CH₂)₂(CH₂)_tR^{1b}, S(CH₂)₂(CH₂)_tR^{1b}, 5-6
 membered carbocycle substituted with 0-1 R⁵, and 5-6
 25 membered heterocycle consisting of: carbon atoms and
 1-4 heteroatoms selected from the group consisting of
 N, O, and S(O)_p substituted with 0-1 R⁵;

R^{4a}, at each occurrence, is selected from H, =O, (CH₂)_rOR²,
 30 (CF₂)_rCF₃, (CH₂)_r-CF₃, (CH₂)_r-F, (CH₂)_r-Br, (CH₂)_r-Cl,

C_{1-4} alkyl, $(CH_2)_rCN$, $(CH_2)_rNO_2$, $(CH_2)_rNR^2R^{2a}$,
 $(CH_2)_rC(O)R^{2c}$, $NR^2C(O)R^{2b}$, $C(O)NR^2R^{2a}$, $(CH_2)_rN=CHOR^3$,
 $C(O)NH(CH_2)_2NR^2R^{2a}$, $NR^2C(O)NR^2R^{2a}$, $C(=NR^2)NR^2R^{2a}$,
 $NHC(=NR^2)NR^2R^{2a}$, $SO_2NR^2R^{2a}$, $NR^2SO_2NR^2R^{2a}$, $NR^2SO_2-C_{1-4}$
5 alkyl, $NR^2SO_2R^5$, $C(O)NHSO_2-C_{1-4}$ alkyl, $S(O)_pR^5$, 5-6
membered carbocycle substituted with 0-1 R^5 , and 5-6
membered heterocycle consisting of: carbon atoms and
1-4 heteroatoms selected from the group consisting of
N, O, and $S(O)_p$, substituted with 0-1 R^5 ;
10
 R^{4b} , at each occurrence, is selected from H, =O, $(CH_2)_rOR^3$,
 $(CH_2)_rF$, $(CH_2)_rCl$, $(CH_2)_rBr$, $(CH_2)_rI$, C_{1-4} alkyl,
 $(CH_2)_rCN$, $(CH_2)_rNO_2$, $(CH_2)_rNR^3R^{3a}$, $(CH_2)_rC(O)R^3$,
 $(CH_2)_rC(O)OR^{3c}$, $NR^3C(O)R^{3a}$, $C(O)NR^3R^{3a}$, $NR^3C(O)NR^3R^{3a}$,
15 $C(=NR^3)NR^3R^{3a}$, $NR^3C(=NR^3)NR^3R^{3a}$, $SO_2NR^3R^{3a}$, $NR^3SO_2NR^3R^{3a}$,
 $NR^3SO_2-C_{1-4}$ alkyl, $NR^3SO_2CF_3$, NR^3SO_2 -phenyl, $S(O)_pCF_3$,
 $S(O)_p-C_{1-4}$ alkyl, $S(O)_p$ -phenyl, $(CH_2)_rCF_3$, and $(CF_2)_rCF_3$;

 R^5 , at each occurrence, is selected from H, C_{1-6} alkyl, =O,
20 $(CH_2)_rOR^3$, F, Cl, Br, I, -CN, NO_2 , $(CH_2)_rNR^3R^{3a}$,
 $(CH_2)_rC(O)R^3$, $(CH_2)_rC(O)OR^{3c}$, $NR^3C(O)R^{3a}$, $C(O)NR^3R^{3a}$,
 $NR^3C(O)NR^3R^{3a}$, $CH(=NOR^{3d})$, $C(=NR^3)NR^3R^{3a}$,
 $NR^3C(=NR^3)NR^3R^{3a}$, $SO_2NR^3R^{3a}$, $NR^3SO_2NR^3R^{3a}$, $NR^3SO_2-C_{1-4}$
alkyl, $NR^3SO_2CF_3$, NR^3SO_2 -phenyl, $S(O)_pCF_3$, $S(O)_p-C_{1-4}$
25 alkyl, $S(O)_p$ -phenyl, $(CF_2)_rCF_3$, phenyl substituted with
0-2 R^6 , naphthyl substituted with 0-2 R^6 , and benzyl
substituted with 0-2 R^6 ;

R⁶, at each occurrence, is selected from H, OH, (CH₂)_rOR², halo, C₁₋₄ alkyl, CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2b}, NR²C(O)R^{2b}, NR²C(O)NR²R^{2a}, C(=NH)NH₂, NHC(=NH)NH₂, SO₂NR²R^{2a}, NR²SO₂NR²R^{2a}, and NR²SO₂C₁₋₄ alkyl;

5

R⁷, at each occurrence, is selected from H, OH, C₁₋₄ alkoxy, C₆₋₁₀ aryloxy, C₆₋₁₀ aryloxyalkyl, C₆₋₁₀ arylmethoxycarbonyl, C₁₋₄ alkylcarbonyloxy C₁₋₄ alkoxy, C₆₋₁₀ arylcarbonyloxy C₁₋₄ alkoxy, C₁₋₆ alkylaminocarbonyl, phenylaminocarbonyl, and phenyl C₁₋₄ alkoxy; and

R⁸, at each occurrence, is selected from H, C₁₋₆ alkyl, and (CH₂)_n-phenyl;

15

alternatively, R⁷ and R⁸, when attached to the same nitrogen, combine to form a 5-6 membered heterocyclic ring consisting of carbon atoms and 0-2 additional heteroatoms selected from the group consisting of N, O, and S(O)_n;

R⁹, at each occurrence, is selected from H, C₁₋₆ alkyl and (CH₂)_n-phenyl;

25 R¹⁰ is selected from H, phenyl substituted with 0-2 R^{4a}, and
naphthyl substituted with 0-2 R^{4a};

n, at each occurrence, is selected from 0, 1, 2, and 3;

30 m. at each occurrence, is selected from 0, 1, and 2;


p, at each occurrence, is selected from 0, 1, and 2;



r, at each occurrence, is selected from 0, 1, 2, and 3;

5 s, at each occurrence, is selected from 0, 1, and 2; and,

t, at each occurrence, is selected from 0, 1, 2, and 3.


10 [2] Thus, in another embodiment, the present invention provides a novel compound selected from the group:





5 or a stereoisomer or pharmaceutically acceptable salt thereof, wherein;

G is selected from formulas Ia₁-Ic₁:



ring D₂ is a 5-membered heteroaromatic ring system
comprising E, carbon atoms, and 0-2 N atoms, wherein E
is selected from O, S, and N-D_{1c} and ring D₂ is
5 substituted with 1 D_{1a} and 0-1 D_{1b};

ring D₃ is a 5-membered heteroaromatic ring system
comprising carbon atoms and from 0-3 additional N atoms
and ring D₃ is substituted with 1 D_{1a} and 0-1 D_{1b};

10 G¹ is selected from H, Cl, F, Br, I, OH, C₁₋₃ alkoxy, NH₂,
NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂, CH₂NH(C₁₋₃
alkyl), CH₂N(C₁₋₃ alkyl)₂, CH₂CH₂NH₂, CH₂CH₂NH(C₁₋₃
alkyl), and CH₂CH₂N(C₁₋₃ alkyl)₂;

15 D_{1a} is selected from H, OH, SH, C₁₋₃ alkoxy, C₁₋₃ thioalkoxy,
NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂, CH₂NH(C₁₋₃
alkyl), CH₂N(C₁₋₃ alkyl)₂, CH₂CH₂NH₂, CH₂CH₂NH(C₁₋₃
alkyl), and CH₂CH₂N(C₁₋₃ alkyl)₂;

20 D_{1b} is selected from H, C₁₋₄ alkyl, Cl, F, Br, I, OH, C₁₋₃
alkoxy, NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂,
CH₂NH(C₁₋₃ alkyl), CH₂N(C₁₋₃ alkyl)₂, CH₂CH₂NH₂,
CH₂CH₂NH(C₁₋₃ alkyl), and CH₂CH₂N(C₁₋₃ alkyl)₂;

25 D_{1c} is selected from H, C₁₋₄ alkyl, C₁₋₃ alkoxy, NH₂, NH(C₁₋₃
alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl),
CH₂N(C₁₋₃ alkyl)₂, CH₂CH₂NH₂, CH₂CH₂NH(C₁₋₃ alkyl), and
CH₂CH₂N(C₁₋₃ alkyl)₂;

Z is selected from a bond, CH_2O , OCH_2 , CH_2NH , NHCH_2 ,
 $\text{NHC(=CH}_2\text{)}$, C(O) , $\text{CH}_2\text{C(O)}$, C(O)CH_2 , NHC(O) , C(O)NH ,
 NHC(O)NH , $\text{CH}_2\text{S(O)}_2$, $\text{S(O)}_2(\text{CH}_2)$, SO_2NH , and NHSO_2 ,
provided that Z does not form a N-N, N-O, NCH_2N , or
5 NCH_2O bond with ring M or group A;

A is selected from one of the following carbocyclic and
heterocyclic systems which are substituted with 0-2 R^4 ;
phenyl, piperidinyl, piperazinyl, pyridyl,
10 pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl,
pyrrolidinyl, oxazolyl, isoxazolyl, thiazolyl,
isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl,
thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl,
1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl,
15 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl,
1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl,
1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl,
1,2,5-triazolyl, 1,3,4-triazolyl, benzofuranyl,
benzothiofuranyl, indolyl, benzimidazolyl,
20 benzoxazolyl, benzthiazolyl, indazolyl, benzisoxazolyl,
benzisothiazolyl, and isoindazolyl;


X is selected from C_{1-4} alkylene, $-\text{C(O)}-$, $-\text{C(=NR)}-$,
 $-\text{CR}^2(\text{NR}^2\text{R}^{2a})-$, $-\text{C(O)CR}^2\text{R}^{2a}-$, $-\text{CR}^2\text{R}^{2a}\text{C(O)}$, $-\text{C(O)NR}^2-$,
25 $-\text{NR}^2\text{C(O)}-$, $-\text{C(O)NR}^2\text{CR}^2\text{R}^{2a}-$, $-\text{NR}^2\text{C(O)CR}^2\text{R}^{2a}-$,
 $-\text{CR}^2\text{R}^{2a}\text{C(O)NR}^2-$, $-\text{CR}^2\text{R}^{2a}\text{NR}^2\text{C(O)}-$, $-\text{NR}^2\text{C(O)NR}^2-$, $-\text{NR}^2-$,
 $-\text{NR}^2\text{CR}^2\text{R}^{2a}-$, $-\text{CR}^2\text{R}^{2a}\text{NR}^2-$, O, $-\text{CR}^2\text{R}^{2a}\text{O}-$, and $-\text{OCR}^2\text{R}^{2a}-$;

alternatively, Y is selected from one of the following
30 carbocyclic and heterocyclic systems which are
substituted with 0-2 R^{4a} ;

cyclopropyl, cyclopentyl, cyclohexyl, phenyl,
 piperidiny, piperaziny, pyridyl, pyrimidyl, furanyl,
 morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl,
 oxazolyl, isoxazolyl, isoxazolinyl, thiazolyl,
 5 isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl,
 thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl,
 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl,
 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl,
 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl,
 10 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl,
 1,2,5-triazolyl, 1,3,4-triazolyl, benzofuranyl,
 benzothiofuranyl, indolyl, benzimidazolyl,
 benzoxazolyl, benzthiazolyl, indazolyl, benzisoxazolyl,
 benzisothiazolyl, and isoindazolyl;

15

alternatively, Y is selected from the following bicyclic
 heteroaryl ring systems:



20 K is selected from O, S, NH, and N;


R^4 , at each occurrence, is selected from H, =O, $(CH_2)_rOR^2$, F,
 Cl, Br, I, C_{1-4} alkyl, CN, NO_2 , $(CH_2)_rNR^2R^{2a}$, $C(O)R^{2c}$,
 $NR^2C(O)R^{2b}$, $C(O)NR^2R^{2a}$, $NR^2C(O)NR^2R^{2a}$, $C(=NR^2)NR^2R^{2a}$,
 25 $SO_2NR^2R^{2a}$, $NR^2SO_2NR^2R^{2a}$, $NR^2SO_2-C_{1-4}$ alkyl, $NR^2SO_2R^5$,
 $S(O)_pR^5$, CF_3 , NCH_2R^{1c} , OCH_2R^{1c} , SCH_2R^{1c} , $N(CH_2)_2(CH_2)_tR^{1b}$,

O(CH₂)₂(CH₂)_tR^{1b}, S(CH₂)₂(CH₂)_tR^{1b}, 5-6 membered carbocycle substituted with 0-1 R⁵, and 5-6 membered heterocycle consisting of: carbon atoms and 1-4 heteroatoms selected from the group consisting of N, O, and S(O)_p, substituted with 0-1 R⁵; and,

R^{4a}, at each occurrence, is selected from H, =O, (CH₂)_rOR², CF₃, F, Br, Cl, C₁₋₄ alkyl, CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2c}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, NR²C(O)NR²R^{2a}, C(=NR²)NR²R^{2a}, NHC(=NR²)NR²R^{2a}, SO₂NR²R^{2a}, NR²SO₂NR²R^{2a}, NR²SO₂-C₁₋₄ alkyl, NR²SO₂R⁵, C(O)NHSO₂-C₁₋₄ alkyl, S(O)_pR⁵, 5-6 membered carbocycle substituted with 0-1 R⁵, and 5-6 membered heterocycle consisting of: carbon atoms and 1-4 heteroatoms selected from the group consisting of N, O, and S(O)_p, substituted with 0-1 R⁵.

[3] Thus, in another embodiment, the present invention provides a novel compound, wherein:

G is selected from formulas Ib₁ and Ic₁:



ring D₂ is a 5-membered heteroaromatic ring system comprising E, carbon atoms, and 0-2 N atoms, wherein E is selected from O, S, and N-D_{1c} and ring D₂ is substituted with 1 D_{1a} and 0-1 D_{1b};

G¹ is selected from H, Cl, F, Br, I, OH, C₁₋₃ alkoxy, NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl), and CH₂N(C₁₋₃ alkyl)₂;

5

D_{1a} is selected from H, OH, SH, NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl), and CH₂N(C₁₋₃ alkyl)₂;

10 D_{1b} is selected from H, C₁₋₄ alkyl, Cl, F, Br, I, OH, C₁₋₃ alkoxy, NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl), and CH₂N(C₁₋₃ alkyl)₂;

D_{1c} is selected from H, C₁₋₄ alkyl, C₁₋₃ alkoxy, NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl), and CH₂N(C₁₋₃ alkyl)₂;

15

Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R^{4a};

20 phenyl, piperidinyl, piperazinyl, pyridyl, pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, imidazolyl, oxadiazole, thiadiazole, triazole, 1,2,3-oxadiazole, 1,2,4-oxadiazole, 1,2,5-oxadiazole, 1,3,4-oxadiazole, 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, 1,3,4-thiadiazole, 1,2,3-triazole, 1,2,4-triazole, 1,2,5-triazole, 1,3,4-triazole, benzofuran, benzothiofuran, indole, benzimidazole, benzimidazolone, benzoxazole, benzthiazole, indazole, benzisoxazole, 30 benzisothiazole, and isoindazole;

Z is selected from a bond, CH_2O , OCH_2 , NH , CH_2NH , NHCH_2 ,
 $\text{CH}_2\text{C}(\text{O})$, $\text{C}(\text{O})\text{CH}_2$, $\text{C}(\text{O})\text{NH}$, $\text{NHC}(\text{O})$, $\text{CH}_2\text{S}(\text{O})_2$, $\text{S}(\text{O})_2(\text{CH}_2)$,
 SO_2NH , and NHSO_2 , provided that Z does not form a N-N,
 5 N-O, N-S, NCH_2N , NCH_2O , or NCH_2S bond with either group
 to which it is attached;


R^4 , at each occurrence, is selected from H, =O, $(\text{CH}_2)_r\text{OR}^2$, F,
 Cl, Br, I, C_{1-4} alkyl, CN, NO_2 , $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$, $\text{C}(\text{O})\text{R}^{2c}$,
 10 $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$, $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$, $\text{NR}^2\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$, $\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$,
 $\text{SO}_2\text{NR}^2\text{R}^{2a}$, $\text{NR}^2\text{SO}_2\text{NR}^2\text{R}^{2a}$, $\text{NR}^2\text{SO}_2\text{-C}_{1-4}$ alkyl, $\text{NR}^2\text{SO}_2\text{R}^5$,
 $\text{S}(\text{O})_p\text{R}^5$, CF_3 , 5-6 membered carbocycle substituted with
 0-1 R^5 , and 5-6 membered heterocycle consisting of:
 carbon atoms and 1-4 heteroatoms selected from the
 15 group consisting of N, O, and $\text{S}(\text{O})_p$ substituted with 0-1
 R^5 ; and,

R^{4a} , at each occurrence, is selected from H, =O, $(\text{CH}_2)_r\text{OR}^2$,
 CF_3 , F, Br, Cl, C_{1-4} alkyl, CN, NO_2 , $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$,
 20 $(\text{CH}_2)_r\text{C}(\text{O})\text{R}^{2c}$, $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$, $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$, $\text{NR}^2\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$,
 $\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$, $\text{SO}_2\text{NR}^2\text{R}^{2a}$, $\text{C}(\text{O})\text{NHSO}_2\text{-C}_{1-4}$ alkyl, $\text{S}(\text{O})_p\text{R}^5$,
 5-6 membered carbocycle substituted with 0-1 R^5 , and 5-6
 membered heterocycle consisting of: carbon atoms and
 1-4 heteroatoms selected from the group consisting of
 25 N, O, and $\text{S}(\text{O})_p$ substituted with 0-1 R^5 .

[4] In a preferred embodiment, the present invention
 provides a novel compound, wherein:

30

G is of formula Ib₂:





- 5 ring D₂ is a 5-membered heteroaromatic ring system comprising E, carbon atoms, and 0-2 N atoms, wherein E is selected from O, S, and N-D_{1c} and ring D₂ is substituted with 1 D_{1a} and 0-1 D_{1b};
- 10 G¹ is selected from H, Cl, F, Br, I, OH, C₁₋₃ alkoxy, NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl), and CH₂N(C₁₋₃ alkyl)₂;
- 15 D_{1a} is selected from H, OH, SH, NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl), and CH₂N(C₁₋₃ alkyl)₂;
- 20 D_{1b} is selected from H, C₁₋₄ alkyl, Cl, F, Br, I, OH, C₁₋₃ alkoxy, NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl), and CH₂N(C₁₋₃ alkyl)₂;
- 25 D_{1c} is selected from H, C₁₋₄ alkyl, C₁₋₃ alkoxy, NH₂, NH(C₁₋₃ alkyl), N(C₁₋₃ alkyl)₂, CH₂NH₂, CH₂NH(C₁₋₃ alkyl), and CH₂N(C₁₋₃ alkyl)₂; and,
- R is selected from H, Cl, F, Br, I, (CH₂)_tOR³, C₁₋₄ alkyl, OCF₃, CF₃, C(O)NR⁷R⁸, (CR⁸R⁹)_tNR⁷R⁸ and (CH₂)_tNR²SO₂-CH₃.

[5] In a more preferred embodiment, the present invention provides a novel compound, wherein:

G is selected from the group:

77





5 Z is selected from $C(O)CH_2$ and $C(O)NH$, provided that Z does not form a N-N bond with group A;

A is selected from phenyl, piperidinyl, pyridyl, and pyrimidyl, and is substituted with 0-2 R^4 ; and,

10

B is selected from phenyl, pyrrolidino, N-pyrrolidino-carbonyl, morpholino, N-morpholino-carbonyl, 1,2,3-triazolyl, imidazolyl, and benzimidazolyl, and is substituted with 0-1 R^{4a} ;

15

R^2 , at each occurrence, is selected from H, CH_3 , CH_2CH_3 , cyclopropylmethyl, cyclobutyl, and cyclopentyl;

R^{2a}, at each occurrence, is selected from H, CH₃, and CH₂CH₃;

alternatively, R² and R^{2a}, together with the atom to which

they are attached, combine to form pyrrolidine

5 substituted with 0-2 R^{4b} or piperidine substituted with
0-2 R^{4b};

R⁴, at each occurrence, is selected from OH, OR², (CH₂)OR²,

(CH₂)₂OR², F, Br, Cl, I, C₁₋₄ alkyl, NR²R^{2a}, (CH₂)NR²R^{2a},

10 (CH₂)₂NR²R^{2a}, CF₃, and (CF₂)CF₃;

R^{4a} is selected from C₁₋₄ alkyl, CF₃, OR², (CH₂)OR²,

(CH₂)₂OR², NR²R^{2a}, (CH₂)NR²R^{2a}, (CH₂)₂NR²R^{2a}, SR⁵, S(O)R⁵,

S(O)₂R⁵, SO₂NR²R^{2a}, and 1-CF₃-tetrazol-2-yl;

15

R^{4b}, at each occurrence, is selected from H, CH₃, and OH;

R⁵, at each occurrence, is selected from CF₃, C₁₋₆ alkyl,


phenyl, and benzyl; and,

20

r, at each occurrence, is selected from 0, 1, and 2.

[6] In an even further preferred embodiment, the present
25 invention provides a novel compound, wherein:

G is selected from:




A is selected from the group: phenyl, piperidinyl, 2-
 5 pyridyl, 3-pyridyl, 2-pyrimidyl, 2-Cl-phenyl, 3-Cl-phenyl, 2-F-phenyl, 3-F-phenyl, 2-methylphenyl, 2-aminophenyl, and 2-methoxyphenyl; and,

B is selected from the group: 2-(aminosulfonyl)phenyl, 2-
 10 (methylaminosulfonyl)phenyl, 1-pyrrolidinocarbonyl, 2-(methylsulfonyl)phenyl, 2-(N,N-dimethylaminomethyl)phenyl, 2-(N-methylaminomethyl)phenyl, 2-(N-ethyl-N-methylaminomethyl)phenyl, 2-(N-pyrrolidinylmethyl)phenyl, 1-methyl-2-imidazolyl, 2-
 15 methyl-1-imidazolyl, 2-(dimethylaminomethyl)-1-imidazolyl, 2-(methylaminomethyl)-1-imidazolyl, 2-(N-(cyclopropylmethyl)aminomethyl)phenyl, 2-(N-(cyclobutyl)aminomethyl)phenyl, 2-(N-

(cyclopentyl)aminomethyl)phenyl, 2-(N-(4-hydroxypiperidinyl)methyl)phenyl, and 2-(N-(3-hydroxypyrrolidinyl)methyl)phenyl.

5

[7] In another even more preferred embodiment, the present invention provides a compound of formula:



10

L_n is $^*CH_2NHC(O)CH_2$ or $^*CH(R^a)NHC(O)CH_2$, the * indicates where L_n is bonded to G;

R^a is $C(O)C(O)OR^3$;

15

Z is selected from a C_{1-4} alkylene, $(CH_2)_rC(O)$, and $(CH_2)_rS(O)_2$;

R^2 , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

20

R^{2a} , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

R^{2b} , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

25

R^{2c} , at each occurrence, is selected from OH, OCH_3 , OCH_2CH_3 , CH_3 , benzyl, and phenyl;

R^3 , at each occurrence, is selected from H, C_{1-4} alkyl, and phenyl;

5 A is C_{5-6} carbocyclic residue substituted with 0-2 R^4 ;

R^4 , at each occurrence, is selected from H, =O, $(CH_2)_rOR^2$, F, Cl, Br, I, C_{1-4} alkyl, -CN, NO_2 , $(CH_2)_rNR^2R^{2a}$, $(CH_2)_rC(O)R^{2c}$, $NR^2C(O)R^{2b}$, $C(O)NR^2R^{2a}$, $C(=NR^2)NR^2R^{2a}$,
10 $NHC(=NR^2)NR^2R^{2a}$, $SO_2NR^2R^{2a}$, $S(O)_pR^5$, and CF_3 ;

R^5 , at each occurrence, is selected from CF_3 , C_{1-6} alkyl, phenyl, and benzyl;

15 p, at each occurrence, is selected from 0, 1, and 2; and,

r, at each occurrence, is selected from 0, 1, 2, and 3.

20 [8] In another still more preferred embodiment, the present invention provides a compound wherein:

L_n is $*CH(R^a)NHC(O)CH_2$;

25 R^a is $C(O)C(O)OH$;

Z is selected from a CH_2 , $(CH_2)_2C(O)$, and $CH_2S(O)_2$;

A is cyclohexyl or phenyl and is substituted with 0-1 R^4 ;


30

R^4 , at each occurrence, is selected from H, =O, OR^2 , CH_2OR^2 ,
 F, Cl, Br, I, C_{1-4} alkyl, -CN, NO_2 , $(CH_2)_rNR^2R^{2a}$,
 $(CH_2)_rC(O)R^{2c}$, $C(O)NR^2R^{2a}$, $SO_2NR^2R^{2a}$, and CF_3 ; and,

5 r , at each occurrence, is selected from 0, 1, and 2.

[9] In another even more preferred embodiment, the present invention provides a compound of formula:

10



L_n is $*CH_2NHC(O)CH_2$ or $*CH(R^a)NHC(O)CH_2$, the * indicates
 where L_n is bonded to G;

15

R^a is $C(O)C(O)OR^3$;

R is H or NH_2 ;

20 Z is selected from a C_{1-4} alkylene, $(CH_2)_rC(O)$, and
 $(CH_2)_rS(O)_2$;

R^2 , at each occurrence, is selected from H, C_{1-6} alkyl,
 benzyl, and phenyl;

25

R^{2a} , at each occurrence, is selected from H, C_{1-6} alkyl,
 benzyl, and phenyl;

R^{2b} , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

R^{2c} , at each occurrence, is selected from OH, OCH_3 , OCH_2CH_3 ,
5 CH_3 , benzyl, and phenyl;

R^3 , at each occurrence, is selected from H, C_{1-4} alkyl, and phenyl;

10 A is a C_{5-6} carbocyclic residue substituted with 0-2 R^4 ;

R^4 , at each occurrence, is selected from H, =O, $(CH_2)_rOR^2$, F, Cl, Br, I, C_{1-4} alkyl, -CN, NO_2 , $(CH_2)_rNR^2R^{2a}$,
 $(CH_2)_rC(O)R^{2c}$, $NR^2C(O)R^{2b}$, $C(O)NR^2R^{2a}$, $SO_2NR^2R^{2a}$, $S(O)_pR^5$,
15 and CF_3 ;

R^5 , at each occurrence, is selected from CF_3 , C_{1-6} alkyl, phenyl, and benzyl;

20 p, at each occurrence, is selected from 0, 1, and 2; and,

r, at each occurrence, is selected from 0, 1, 2, and 3.

25 [10] In another still more preferred embodiment, the present invention provides a compound wherein:

L_n is $*CH(R^a)NHC(O)CH_2$;

30 R is H;

R^a is $C(O)C(O)OH$;

Z is selected from a CH_2 , $(CH_2)_2C(O)$, and $CH_2S(O)_2$;

5 A is cyclohexyl or phenyl and is substituted with 0-1 R^4 ;

R^4 , at each occurrence, is selected from H , $=O$, OR^2 , CH_2OR^2 ,


F , Cl , Br , I , C_{1-4} alkyl, $-CN$, NO_2 , $(CH_2)_rNR^2R^{2a}$,

$(CH_2)_rC(O)R^{2c}$, $C(O)NR^2R^{2a}$, $SO_2NR^2R^{2a}$, and CF_3 ;

10

r , at each occurrence, is selected from 0, 1, 2, and 3.

[11] In another even more preferred embodiment, the present
15 invention provides a compound of formula:



L_n is $*CH_2NHC(O)CH_2$ or $*CH(R^a)NHC(O)CH_2$, the $*$ indicates
20 where L_n is bonded to G ;

R is H or NH_2 ;

R^a is $C(O)C(O)OR^3$;

25

Z is C_{1-4} alkylene;

R^2 , at each occurrence, is selected from H , C_{1-6} alkyl,
benzyl, and phenyl;

R^{2a}, at each occurrence, is selected from H, C₁₋₆ alkyl, benzyl, and phenyl;

5 R^{2b}, at each occurrence, is selected from H, C₁₋₆ alkyl, benzyl, and phenyl;

R^{2c}, at each occurrence, is selected from OH, OCH₃, OCH₂CH₃, CH₃, benzyl, and phenyl;

10

R³, at each occurrence, is selected from H, C₁₋₄ alkyl, and phenyl;

A is phenyl substituted with 0-2 R⁴;

15

R⁴, at each occurrence, is selected from H, (CH₂)_rOR², F, Cl, Br, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2c}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, S(O)_pR⁵, and CF₃;

20 R⁵, at each occurrence, is selected from CF₃, C₁₋₆ alkyl, phenyl, and benzyl;

p, at each occurrence, is selected from 0, 1, and 2; and,

25 r, at each occurrence, is selected from 0, 1, 2, and 3.

[12] In another still more preferred embodiment, the present invention provides a compound wherein:

30

L_n is *CH(R^a)NHC(O)CH₂;

R is NH_2 ;

R^a is $\text{C}(\text{O})\text{C}(\text{O})\text{OH}$;

5

Z is CH_2 ;


A is phenyl substituted with 0-1 R^4 ;

10 R^4 , at each occurrence, is selected from H, OR^2 , CH_2OR^2 , F, Cl, Br, C_{1-4} alkyl, $-\text{CN}$, NO_2 , $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$, $(\text{CH}_2)_r\text{C}(\text{O})\text{R}^{2c}$, $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$, $\text{SO}_2\text{NR}^2\text{R}^{2a}$, and CF_3 ; and,

r, at each occurrence, is selected from 0, 1, and 2.

15

[13] In another even more preferred embodiment, the present invention provides a compound of formula:



20

L_n is $^*\text{CH}_2\text{NHC}(\text{O})$ or $^*\text{CH}(\text{R}^a)\text{NHC}(\text{O})$ and the * indicates where L_n is bonded to G;

R^a is selected from $\text{C}(\text{O})\text{C}(\text{O})\text{OR}^3$ and $\text{C}(\text{O})-\text{A}$;

25

R^b is selected from H, phenyl, C_{1-10} alkyl, and C_{2-5} alkenyl;

R^c is selected from H and C_{1-6} alkyl;

alternatively, R^b and R^c together are -(CH₂)₄-;

Z is (CR⁸R⁹)₁₋₄;

5

R², at each occurrence, is selected from H, CF₃, and C₁₋₆ alkyl;

10 R^{2a}, at each occurrence, is selected from H, CF₃, and C₁₋₆ alkyl;

R^{2b}, at each occurrence, is selected from H, CF₃, and C₁₋₆ alkyl;

15 R^{2c}, at each occurrence, is selected from OH, OCH₃, OCH₂CH₃, CH₃, benzyl, and phenyl;

R³, at each occurrence, is selected from H, C₁₋₄ alkyl, and phenyl;

20

A is selected from:

C₆₋₁₀ aromatic carbocyclic residue substituted with 0-2 R⁴, and

25 5-10 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R⁴;

30 R⁴, at each occurrence, is selected from H, (CH₂)_rOR², F, Cl, Br, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2c}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, S(O)_pR⁵, and CF₃;

R⁵, at each occurrence, is selected from CF₃, C₁₋₆ alkyl, phenyl, and benzyl;

5 R⁸, at each occurrence, is selected from H, C₁₋₆ alkyl and phenyl;

R⁹, at each occurrence, is selected from H, C₁₋₆ alkyl and phenyl;

10

p, at each occurrence, is selected from 0, 1, and 2;

r, at each occurrence, is selected from 0, 1, 2, and 3.

15

[14] In another still more preferred embodiment, the present invention provides a compound wherein:

20 L_n is *CH(R^a)NHC(O) and the * indicates where L_n is bonded to G;

R^a is C(O)C(O)OH or C(O)-(benzothiazol-2-yl);

25 R^b is selected from H, phenyl, C₁₋₁₀ alkyl, and C₂₋₅ alkenyl;

R^c is selected from H and C₁₋₆ alkyl;

alternatively, R^b and R^c together are -(CH₂)₄-;

30 Z is (CR⁸H)₁₋₂;

A is selected from phenyl, naphthyl, and thienyl, and A is substituted with 0-1 R⁴;


R⁴, at each occurrence, is selected from H, OR², CH₂OR², F,
 5 Cl, Br, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a},
 (CH₂)_rC(O)R^{2c}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, and CF₃;

R⁸, at each occurrence, is selected from H, methyl and
 phenyl; and,

10

r, at each occurrence, is selected from 0, 1, and 2.

[15] In another even more preferred embodiment, the present
 15 invention provides a compound of formula:



L_n is *CH₂NHC(O) or *CH(R^a)NHC(O) and the * indicates where
 L_n is bonded to G;

20

R^a is selected from C(O)C(O)OR³ and C(O)-A;

R^b is selected from H, phenyl, C₁₋₁₀ alkyl, and C₂₋₅ alkenyl;

25 R^c is selected from H and C₁₋₆ alkyl;

alternatively, R^b and R^c together are -(CH₂)₄-;

R is selected from H, benzyl, C₁₋₄ alkyl, and NH₂;

Z is (CR⁸R⁹)₁₋₄;

5 R², at each occurrence, is selected from H, CF₃, and C₁₋₆ alkyl;

R^{2a}, at each occurrence, is selected from H, CF₃, and C₁₋₆ alkyl;

10

R^{2b}, at each occurrence, is selected from H, CF₃, and C₁₋₆ alkyl;

15 R^{2c}, at each occurrence, is selected from OH, OCH₃, OCH₂CH₃, CH₃, benzyl, and phenyl;

R³, at each occurrence, is selected from H, C₁₋₄ alkyl, and phenyl;

20 A is selected from:

C₆₋₁₀ aromatic ring substituted with 0-2 R⁴, and
5-10 membered aromatic heterocyclic system containing
from 1-4 heteroatoms selected from the group consisting of
N, O, and S substituted with 0-2 R⁴;

25

R⁴, at each occurrence, is selected from H, (CH₂)_rOR², F, Cl, Br, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2c}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, S(O)_pR⁵, and CF₃;

R⁵, at each occurrence, is selected from CF₃, C₁₋₆ alkyl, phenyl, and benzyl;

5 R⁸, at each occurrence, is selected from H, C₁₋₆ alkyl and phenyl;

R⁹, at each occurrence, is selected from H, C₁₋₆ alkyl and phenyl;

10 p, at each occurrence, is selected from 0, 1, and 2;

r, at each occurrence, is selected from 0, 1, 2, and 3.

15 [16] In another still more preferred embodiment, the present invention provides a compound wherein:

L_n is *CH(R^a)NHC(O) and the * indicates where L_n is bonded to G;

20

R^a is C(O)C(O)OH or C(O)-(benzothiazol-2-yl);

R^b is selected from H, phenyl, C₁₋₁₀ alkyl, and C₂₋₅ alkenyl;

25 R^c is selected from H and C₁₋₆ alkyl;

alternatively, R^b and R^c together are -(CH₂)₄-;

Z is (CR⁸H)₁₋₂;

30

A is selected from phenyl, naphthyl, and thienyl, and A is substituted with 0-1 R⁴;


R⁴, at each occurrence, is selected from H, OR², CH₂OR², F,
 5 Cl, Br, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a},
 (CH₂)_rC(O)R^{2c}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, and CF₃;

R⁸, at each occurrence, is selected from H, C₁₋₆ alkyl and
 phenyl;

10

r, at each occurrence, is selected from 0, 1, and 2.

[17] In another even more preferred embodiment, the present
 15 invention provides a compound of formula:



L_n is *CH₂NHC(O) or *CH(R^a)NHC(O) and the * indicates where
 L_n is bonded to G;

20

R^{1a} is selected from -(CH₂)_r-R^{1b} and NHCH₂R^{1c};

R^{1b} is selected from H, OR², NR²R^{2a}, and NR²SO₂(CH₂)_rR^{2b};

25 R^{1c} is selected from C(O)NR²R^{2a}, S(O)₂R^{2b}, and SO₂NR²R^{2a};

R², at each occurrence, is selected from H, C₁₋₆ alkyl,
 benzyl, and phenyl;

R^{2a}, at each occurrence, is selected from H, C₁₋₆ alkyl, benzyl, and phenyl;

5 R^{2b}, at each occurrence, is selected from C₁₋₄ alkoxy, C₁₋₆ alkyl, benzyl, phenyl substituted with 0-2 R^{4b}, and 5-6 membered heterocyclic system containing from 1-2 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4b};

10 R^{2c}, at each occurrence, is selected from OH, OCH₃, OCH₂CH₃, CH₃, benzyl, and phenyl;

alternatively, R² and R^{2a}, together with the atom to which they are attached, combine to form a 5 or 6 membered
15 saturated, partially saturated or unsaturated ring substituted with 0-2 R^{4b} and containing from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

20 R³, at each occurrence, is selected from H, C₁₋₄ alkyl, and phenyl;

A is phenyl substituted with 0-2 R⁴;

25 A¹ is H or A;

alternatively, A and A¹ and the carbon to which they are attached combine to form fluorene;

30 A² is selected from H, A, and CHA³A⁴;

A³ is selected from H, A, C₁₋₄ alkyl, and -(CH₂)_rNR²R^{2a};

A⁴ is H or A;

5 R⁴, at each occurrence, is selected from H, (CH₂)_rOR², F, Cl, Br, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2c}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, S(O)_pR⁵, and CF₃;

10 R^{4b}, at each occurrence, is selected from H, (CH₂)_rOR², F, Cl, Br, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2c}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, S(O)_pR⁵, and CF₃;

15 R⁵, at each occurrence, is selected from CF₃, C₁₋₆ alkyl, phenyl, and benzyl;

p, at each occurrence, is selected from 0, 1, and 2;

20 r, at each occurrence, is selected from 0, 1, 2, and 3.

[18] In another still more preferred embodiment, the present invention provides a compound wherein:

25 L_n is *CH₂NHC(O) and the * indicates where L_n is bonded to G;

R^{1a} is selected from -(CH₂)_r-R^{1b} and NHCH₂R^{1c};

30 R^{1b} is selected from OH, NR²R^{2a}, and NR²SO₂(CH₂)_rR^{2b};

R^{1c} is selected from C(O)NR²R^{2a}, S(O)₂R^{2b}, and SO₂NR²R^{2a};

R^2 , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

5 R^{2a} , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

R^{2b} , at each occurrence, is selected from C_{1-4} alkoxy, C_{1-6} alkyl, benzyl, phenyl substituted with 0-1 R^{4b} , and
10 pyrrolidinyl substituted with 0-1 R^{4b} ;

R^{2c} , at each occurrence, is selected from OH, OCH_3 , OCH_2CH_3 , CH_3 , benzyl, and phenyl;

15 alternatively, R^2 and R^{2a} , together with the atom to which they are attached, combine to form a piperidine ring substituted with 0-1 R^{4b} ;


R^4 , at each occurrence, is selected from H, =O, OR^2 , CH_2OR^2 ,
20 F, Cl, Br, I, C_{1-4} alkyl, -CN, NO_2 , $(CH_2)_rNR^2R^{2a}$,
 $(CH_2)_rC(O)R^{2c}$, $C(O)NR^2R^{2a}$, $SO_2NR^2R^{2a}$, and CF_3 ;

R^{4b} , at each occurrence, is selected from H, =O, OH, F, Cl, C_{1-4} alkyl, and NH_2 ; and,

25

r , at each occurrence, is selected from 0, 1, and 2.

[19] In another even more preferred embodiment, the present
30 invention provides a compound of formula:



L_n is CH_2 ;

R^{1a} is $-(\text{CH}_2)_r-R^{1b}$;

5

R^{1b} is selected from H, C_{1-3} alkyl, $(\text{CH}_2)_r\text{OR}^2$, NR^2R^{2a} , $\text{C}(\text{O})\text{R}^{2c}$, phenyl substituted with 0-2 R^4 , and 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^4 ;

10

R^2 , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

15 R^{2a} , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

R^{2b} , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

20

R^{2c} , at each occurrence, is selected from OH, OCH_3 , OCH_2CH_3 , CH_3 , benzyl, and phenyl;

R^3 , at each occurrence, is selected from H, C_{1-4} alkyl, and phenyl;

25

A is selected from:

C_6-10 aromatic ring substituted with 0-2 R^4 , and

5-10 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^4 ;

5 B is selected from: H, Y, and X-Y

X is selected from C_{1-4} alkylene, $-NR^2-$, and O;

Y is selected from:

10 C_{6-10} aromatic ring substituted with 0-2 R^{4a} , and
5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4a} ;

15 R^4 , at each occurrence, is selected from H, $(CH_2)_rOR^2$, F, Cl, Br, I, C_{1-4} alkyl, $-CN$, NO_2 , $(CH_2)_rNR^2R^{2a}$, $(CH_2)_rC(O)R^{2c}$, $NR^2C(O)R^{2b}$, $C(O)NR^2R^{2a}$, $SO_2NR^2R^{2a}$, $S(O)_pR^5$, and CF_3 ;

R^{4a} , at each occurrence, is selected from H, $(CH_2)_rOR^2$, Cl, Br, F, I, C_{1-4} alkyl, $-CN$, NO_2 , $(CH_2)_rNR^2R^{2a}$, $(CH_2)_rC(O)R^{2c}$, $NR^2C(O)R^{2b}$, $C(O)NR^2R^{2a}$, $SO_2NR^2R^{2a}$, $S(O)_pR^5$, and CF_3 ;

20

R^5 , at each occurrence, is selected from CF_3 , C_{1-6} alkyl, phenyl, and benzyl;

25

p, at each occurrence, is selected from 0, 1, and 2; and,

r, at each occurrence, is selected from 0, 1, 2, and 3.

30

[20] In another still more preferred embodiment, the present invention provides a compound wherein:

R^{1a} is $-(CH_2)_r-R^{1b}$;

5

R^{1b} is selected from H, C₁₋₃ alkyl, OH, NR²R^{2a}, and phenyl substituted with 0-2 R⁴;

A is selected from:

10 phenyl substituted with 0-2 R⁴, naphthyl substituted with 0-2 R⁴, thienyl substituted with 0-2 R⁴, benzothienyl substituted with 0-2 R⁴, 5-aza-benzothienyl substituted with 0-2 R⁴, 6-azabenzothienyl substituted with 0-2 R⁴, and quinolinyl substituted with 0-2 R⁴;

15

B is selected from: H, Y, and X-Y

X is O;

20 Y is phenyl substituted with 0-1 R^{4a};

R⁴, at each occurrence, is selected from H, OR², CH₂OR², F, Cl, Br, I, C₁₋₄ alkyl, -CN, (CH₂)_rNR²R^{2a}, C(O)NR²R^{2a}, and CF₃;

25

R^{4a}, at each occurrence, is selected from H, OR², CH₂OR², F, Cl, Br, I, C₁₋₄ alkyl, -CN, (CH₂)_rNR²R^{2a}, C(O)NR²R^{2a}, and CF₃; and,

30 r, at each occurrence, is selected from 0, 1, and 2.

[21] In another even more preferred embodiment, the present invention provides a compound of formula:



5

L_n is O or S;

M^2 is N or CR^f ;

10 M^3 is N or CR^d ;

provided that only one of M^2 and M^3 is N;

15 R^e is selected from H, $N(CH_3)(CH_2CO_2H)$ and S-(5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^4);

R^d is selected from H, F, and Cl;

20

alternatively, R^d and R^e combine to form $-NR^3-C(O)-C(R^{1g}R^3)-NR^3-$ or $-N=CR^2-NR^3-$;

R^f is selected from H, F, and Cl;

25

alternatively, R^e and R^f combine to form $-NR^3-C(R^{1g}R^3)-C(O)-NR^3-$ or $-NR^3-CR^2=N-$;

Z is O, provided that Z does not form a N-O or NCH₂O bond
with the groups to which Z is attached;

5 R^{1g} is selected from H, C₁₋₆ alkyl, and C₁₋₆ alkyl substituted
with A;

R², at each occurrence, is selected from H, C₁₋₆ alkyl,
benzyl, and phenyl;

10 R^{2a}, at each occurrence, is selected from H, C₁₋₆ alkyl,
benzyl, and phenyl;

R^{2b}, at each occurrence, is selected from H, C₁₋₆ alkyl,
benzyl, and phenyl;

15 R^{2c}, at each occurrence, is selected from OH, OCH₃, OCH₂CH₃,
CH₃, benzyl, and phenyl;

20 R³, at each occurrence, is selected from H, C₁₋₄ alkyl, and
phenyl;

A is selected from:

C₅₋₆ carbocyclic residue substituted with 0-2 R⁴, and
5-6 membered heterocyclic system containing from 1-4
25 heteroatoms selected from the group consisting of N, O, and
S substituted with 0-2 R⁴;

B is H or Y;

30 Y is selected from:

C₅₋₆ carbocyclic residue substituted with 0-2 R^{4a}, and

5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R^{4a};

5 R⁴, at each occurrence, is selected from H, =O, (CH₂)_rOR², F, Cl, Br, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2c}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, C(=NR²)NR²R^{2a}, NHC(=NR²)NR²R^{2a}, SO₂NR²R^{2a}, and CF₃;

10 R^{4a}, at each occurrence, is selected from H, =O, (CH₂)_rOR², (CH₂)_r-F, (CH₂)_r-Br, (CH₂)_r-Cl, Cl, Br, F, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2c}, NR²C(O)R^{2b}, C(O)NR²R^{2a}, C(=NR²)NR²R^{2a}, NHC(=NR²)NR²R^{2a}, SO₂NR²R^{2a}, and CF₃; and,

15

r, at each occurrence, is selected from 0, 1, 2, and 3.

[22] In another still more preferred embodiment, the present
20 invention provides a compound wherein:

L_n is O;

R^e is N(CH₃)(CH₂CO₂H);

25

R^d is H or F;

alternatively, R^d and R^e combine to form -NR³-C(O)-C(R^{1g}R³)-
NR³- or -N=CR²-NR³-;

30

R^f is H or F;

alternatively, R^e and R^f combine to form $-NR^3-C(R^{1g}R^3)-C(O)-$
 NR^3- or $-NR^3-CR^2=N-$;

5

R^{1g} is selected from H, C_{1-2} alkyl and benzyl;

A is phenyl substituted with 0-2 R^4 ;

10 B is H or Y;

Y is 5 membered heterocyclic system containing from 1-2
heteroatoms selected from the group consisting of N, O,
and S substituted with 0-2 R^{4a} ;

15


R^4 , at each occurrence, is selected from H, C_{1-4} alkyl, and
 NR^2R^{2a} ; and,

R^{4a} , at each occurrence, is selected from H, C_{1-4} alkyl, and
 NR^2R^{2a} .

20

[23] In another even more preferred embodiment, the present
invention provides a compound of formula:

25



L_n is $*CH_2NHC(O)CH_2$ or $*CH(R^a)NHC(O)CH_2$ and the * indicates
where L_n is bonded to G;

R^a is C(O)C(O)OR³;

R, at each occurrence, is selected from H, Cl, F, Br, I,
OR³, C₁₋₄ alkyl, C(O)NH₂, and NH₂;

5

Z is selected from a C₁₋₄ alkylene and (CH₂)_rSO₂NR³;

R², at each occurrence, is selected from H, C₁₋₆ alkyl,
benzyl, and phenyl;

10

R^{2a}, at each occurrence, is selected from H, C₁₋₆ alkyl,
benzyl, and phenyl;

R^{2c}, at each occurrence, is selected from OH, OCH₃, OCH₂CH₃,
CH₃, benzyl, and phenyl;

15

R³, at each occurrence, is selected from H, C₁₋₄ alkyl, and
phenyl;

20 A is selected from:

C₅₋₆ carbocyclic residue substituted with 0-2 R⁴, and
5-6 membered aromatic heterocyclic system containing
from 1-4 heteroatoms selected from the group consisting of
N, O, and S substituted with 0-2 R⁴;

25

B is selected from: H, Y, and X-Y

alternatively, when B is H, A is (phenyl)₂CH- substituted
with 0-2 R⁴;

30

X is selected from C₁₋₄ alkylene, -C(O)-, -NR²-, and O;

Y is selected from:

C₅₋₆ carbocyclic residue substituted with 0-2 R^{4a}, and
5 5-6 membered aromatic heterocyclic system containing
from 1-4 heteroatoms selected from the group consisting of
N, O, and S substituted with 0-2 R^{4a};

R⁴, at each occurrence, is selected from H, =O, (CH₂)_rOR², F,
Cl, Br, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a},
10 (CH₂)_rC(O)R^{2c}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, and CF₃;

R^{4a}, at each occurrence, is selected from H, =O, (CH₂)_rOR²,
Cl, Br, F, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a},
(CH₂)_rC(O)R^{2c}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, and CF₃; and,
15

r, at each occurrence, is selected from 0, 1, 2, and 3.

[24] In another still more preferred embodiment, the present
20 invention provides a compound wherein:

L_n is *CH₂NHC(O)CH₂ and the * indicates where L_n is bonded to
G;

25 R, at each occurrence, is selected from H and C₁₋₄ alkyl;

Z is CH₂SO₂NR³;

A is phenyl substituted with 0-2 R⁴;

30


B is H;

R^4 , at each occurrence, is selected from H, $(CH_2)_rOR^2$, F, Cl, C_{1-4} alkyl, $(CH_2)_rNR^2R^{2a}$, $(CH_2)_rC(O)R^{2c}$, and $C(O)NR^2R^{2a}$; and,

5

r , at each occurrence, is selected from 0, 1, and 2.

[25] In another even more preferred embodiment, the present invention provides a compound of formula:



L_n is $*CH_2NHC(O)CH_2$ or $*CH(R^a)NHC(O)CH_2$ and the $*$ indicates where L_n is bonded to G;

15

R^a is $C(O)C(O)OR^3$;

R , at each occurrence, is selected from H, C_{1-4} alkyl, and NH_2 ;

20

R^{1g} is H or C_{1-6} alkyl;

Z is selected from a C_{1-4} alkylene and $(CH_2)_rS(O)_p(CH_2)_r$;

R^2 , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

R^{2a} , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

R^{2c}, at each occurrence, is selected from OH, OCH₃, OCH₂CH₃,
CH₃, benzyl, and phenyl;

5 R³, at each occurrence, is selected from H, C₁₋₄ alkyl, and
phenyl;

A is selected from:

10 C₃₋₆ carbocyclic residue substituted with 0-2 R⁴, and
5-6 membered aromatic heterocyclic system containing
from 1-4 heteroatoms selected from the group consisting of
N, O, and S substituted with 0-2 R⁴;

B is selected from: H, Y, and X-Y

15

alternatively, when B is H, A is (phenyl)₂CH- substituted
with 0-2 R⁴;

X is selected from C₁₋₄ alkylene, -C(O)-, -NR²-, and O;

20

Y is selected from:

C₅₋₆ carbocyclic residue substituted with 0-2 R^{4a}, and
5-6 membered aromatic heterocyclic system containing
from 1-4 heteroatoms selected from the group consisting of
25 N, O, and S substituted with 0-2 R^{4a};

alternatively, Z-A-B combine to form S-C₁₋₆ alkyl;

30 R⁴, at each occurrence, is selected from H, =O, (CH₂)_rOR², F,
Cl, Br, I, C₁₋₄ alkyl, -CN, NO₂, (CH₂)_rNR²R^{2a},
(CH₂)_rC(O)R^{2c}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, and CF₃;

R^{4a} , at each occurrence, is selected from H, =O, $(CH_2)_rOR^2$,
Cl, Br, F, I, C_{1-4} alkyl, -CN, NO_2 , $(CH_2)_rNR^2R^{2a}$,
 $(CH_2)_rC(O)R^{2c}$, $C(O)NR^2R^{2a}$, $SO_2NR^2R^{2a}$, and CF_3 ;

5

p is selected from 0, 1, and 2; and,

r, at each occurrence, is selected from 0, 1, 2, and 3.

10

[26] In another still more preferred embodiment, the present invention provides a compound wherein:

L_n is $*CH_2NHC(O)CH_2$ and the * indicates where L_n is bonded to
15 G;

R is H or C_{1-4} alkyl;

R^{1g} is H;

20

Z is CH_2 , CH_2S , or $CH_2S(O)_2$;

A is a C_{3-6} carbocyclic residue substituted with 0-2 R^4 ;


25 B is H

alternatively, Z-A-B combine to form S- C_{1-6} alkyl;

R^4 , at each occurrence, is selected from H, $(CH_2)_rOR^2$, F, Cl,
30 Br, C_{1-4} alkyl, $(CH_2)_rNR^2R^{2a}$, $(CH_2)_rC(O)R^{2c}$, $C(O)NR^2R^{2a}$,
 $SO_2NR^2R^{2a}$, and CF_3 ; and,

r, at each occurrence, is selected from 0, 1, and 2.

- 5 [27] In another even more preferred embodiment, the present invention provides a compound of formula:



10 L_n is $*CH_2NHC(O)CH_2$ or $*CH(R^a)NHC(O)CH_2$ and the * indicates where L_n is bonded to G;

M^1 is absent or is selected from CHR, O, and NR^2 ;

15 M^4 is selected from NR^2 , CR^f , and $C(O)$;

R is selected from H, Cl, F, Br, I, OR^3 , C_{1-4} alkyl, OCF_3 , CF_3 , and NH_2 ;

20 Z is C_{1-4} alkylene;

R^2 , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

25 R^{2a} , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

R^{2c} , at each occurrence, is selected from OH, OCH_3 , OCH_2CH_3 , CH_3 , benzyl, and phenyl;

R³, at each occurrence, is selected from H, C₁₋₄ alkyl, and phenyl;

A is selected from:

5 C₃₋₆ carbocyclic residue substituted with 0-2 R⁴, and 5-6 membered aromatic heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R⁴;

10 R⁴, at each occurrence, is selected from H, (CH₂)_rOR², F, Cl, Br, I, C₁₋₄ alkyl, (CH₂)_rNR²R^{2a}, (CH₂)_rC(O)R^{2c}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, and CF₃; and,

r, at each occurrence, is selected from 0, 1, 2, and 3.

15

[28] In another still more preferred embodiment, the present invention provides a compound wherein:

20 L_n is *CH₂NHC(O)CH₂ and the * indicates where L_n is bonded to G;

M¹ is absent;

25 R is selected from H and C₁₋₄ alkyl;

Z is CH₂;

A is C₃₋₆ carbocyclic residue substituted with 0-1 R⁴;


30

R^4 , at each occurrence, is selected from H, C_{1-4} alkyl, $(CH_2)_rNR^2R^{2a}$, and CF_3 ; and,

r , at each occurrence, is selected from 0, 1, and 2.

5

[29] In another even more preferred embodiment, the present invention provides a compound of formula:



10

L_n is $*CH_2NHC(O)CH_2$ or $*CH(R^a)NHC(O)CH_2$ and the * indicates where L_n is bonded to G;

R^a is $C(O)C(O)OR^3$;

15

R , at each occurrence, is selected from H, Cl, F, Br, I, OR^3 , C_{1-4} alkyl, $C(O)NH_2$, and NH_2 ;

Z is $(CHR^8)NR^3$, $(CHR^8)_2NR^3$, and $(CHR^8)_2SO_2R^3$;

20

provided that when Z is $(CHR^8)_2NR^3$, then B is absent;

R^2 , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

25

R^{2a} , at each occurrence, is selected from H, C_{1-6} alkyl, benzyl, and phenyl;

R^{2c}, at each occurrence, is selected from OH, OCH₃, OCH₂CH₃,
CH₃, benzyl, and phenyl;

5 R³, at each occurrence, is selected from H, C₁₋₄ alkyl, and
phenyl;

R^{3a}, at each occurrence, is selected from H, C₁₋₄ alkyl, and
phenyl;

10 B is H or Y;

Y is selected from:

C₅₋₆ carbocyclic residue substituted with 0-2 R^{4a}, and
5-6 membered heterocyclic system containing from 1-2
15 heteroatoms selected from the group consisting of N, O, and
S substituted with 0-2 R^{4a};

R^{4a}, at each occurrence, is selected from H, =O, (CH₂)_rOR²,
Cl, Br, F, I, C₁₋₄ alkyl, -CN, (CH₂)_rNR²R^{2a},
20 (CH₂)_rC(O)R^{2c}, C(O)NR²R^{2a}, SO₂NR²R^{2a}, and CF₃;

R⁸, at each occurrence, is selected from H, C₁₋₆ alkyl and
phenyl; and,

25 r, at each occurrence, is selected from 0, 1, 2, and 3.

[30] In another still more preferred embodiment, the present
invention provides a compound wherein:

30

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number
WO 02/002519 A3

(51) International Patent Classification⁷: **C07D 487/04**,
A61K 31/50 // (C07D 487/04, 241:00), 209:00)

(21) International Application Number: PCT/US01/20962

(22) International Filing Date: 29 June 2001 (29.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/214,831 29 June 2000 (29.06.2000) US

(71) Applicant: **BRISTOL-MYERS SQUIBB PHARMA COMPANY** [US/US]; Patent Dept., P.O. Box 4000, Princeton, NJ 08543-4000 (US).

(72) Inventors: **PINTO, Donald, J., P.**; 425 Brentford Road, Kennett Square, PA 19348 (US). **QUAN, Mimi, L.**; 113 Venus Drive, Newark, DE 19711 (US).

(74) Agent: **BELFIELD, Jing, S.**; Bristol-Myers Squibb Pharma Company, Route 141 & Henry Clay Road, Experimental Station, E353/133B, Wilmington, DE 19898-0353 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
25 July 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/002519 A3

(54) Title: THROMBIN OR FACTOR Xa INHIBITORS

(57) Abstract: This invention relates generally to heteroaryl-phenyl substituted compounds that are inhibitors of trypsin-like serine protease enzymes, especially factor Xa or thrombin, pharmaceutical compositions containing the same, and methods of using the same as anticoagulant agents for treatment and prevention of thromboembolic disorders.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/20962

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 A61K31/50 A61K31/50 A61K31/50 A61K31/50
 //(C07D487/04,241:00,209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 28326 A (SIDDIQUI M ARSHAD ;BACHAND BENOIT (CA); IAF BIOCHEM INT (CA); EDMU) 2 July 1998 (1998-07-02) cited in the application claim 28, 2 last cpds; ex. page 40 ---	1-8, 54-57
Y	WO 97 48706 A (SIDDIQUI M ARSHAD ;EDMUNDS JEREMY JOHN (US); WARNER LAMBERT CO (US) 24 December 1997 (1997-12-24) abstract; claims ---	1-8, 54-57
Y	WO 96 19483 A (IAF BIOCHEM INT ;DIMAIO JOHN (CA); SIDDIQUI M ARSHAD (CA); GILLARD) 27 June 1996 (1996-06-27) cited in the application cpds VIII, ex. 495,510,515,640,760,765 --- -/--	1-8, 54-57

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

4 February 2002

Date of mailing of the international search report

16. 5. 02

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Frelon, D

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/20962

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	L.S. NARASIMHAN ET AL.: "Structural basis of the thrombin selectivity..." JOURNAL OF MEDICINAL CHEMISTRY, vol. 43, 2000, pages 361-368, XP002188952 WASHINGTON US the whole document	1-8, 54-57
Y	ST-DENIS Y ET AL: "Potent bicyclic lactam inhibitors of thrombin: Part I: P3 modifications" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 8, no. 22, 17 November 1998 (1998-11-17), pages 3193-3198, XP004143725 ISSN: 0960-894X the whole document	1-8, 54-57
Y	PLUMMER J S ET AL: "Potent and selective bicyclic lactam inhibitors of thrombin: part 2: P1 modifications" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 8, no. 23, 1 December 1998 (1998-12-01), pages 3409-3414, XP004143767 ISSN: 0960-894X the whole document	1-8, 54-57
Y	PLUMMER J S ET AL: "Potent and selective bicyclic lactam inhibitors of thrombin: part 3: P1' modifications" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 9, no. 6, 22 March 1999 (1999-03-22), pages 835-840, XP004160484 ISSN: 0960-894X the whole document	1-8, 54-57
A	WO 98 57937 A (DU PONT MERCK PHARMA) 23 December 1998 (1998-12-23) page 191; example 73	1-8, 54-57
A	WO 97 23212 A (DU PONT MERCK PHARMA) 3 July 1997 (1997-07-03) claim 1: cpds IIIa, IIIb	1-8, 54-57

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/20962

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 53
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-6,7,8,54,56-57 all in part

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-6(in part),7,8,54(in part),56-57(in part)

Compounds first designated in claim 1, corresponding to compounds claimed in claim 7 and dependent claims thereof

2. Claims: 1-6(in part),9,10,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 9 and dependent claims thereof

3. Claims: 1-6(in part),11,12,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 11 and dependent claims thereof

4. Claims: 1-6(in part),13,14,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 13 and dependent claims thereof

5. Claims: 1-6(in part),15,16,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 15 and dependent claims thereof

6. Claims: 1-6(in part),17,18,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 17 and dependent claims thereof

7. Claims: 1-6(in part),19,20,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 19 and dependent claims thereof

8. Claims: 1-6(in part),21,22,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 21 and dependent claims thereof

9. Claims: 1-6(in part),23,24,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 23 and dependent claims thereof

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

10. Claims: 1-6(in part),25,26,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 25 and dependent claims thereof

11. Claims: 1-6(in part),27,28,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 27 and dependent claims thereof

12. Claims: 1-6(in part),29,30,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 29 and dependent claims thereof

13. Claims: 1-6(in part),31,32,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 31 and dependent claims thereof

14. Claims: 1-6(in part),33,34,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 34 and dependent claims thereof

15. Claims: 1-6(in part),35,36,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 35 and dependent claims thereof

16. Claims: 1-6(in part),37,38,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 37 and dependent claims thereof

17. Claims: 1-6(in part),39,40,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 39 and dependent claims thereof

18. Claims: 1-6(in part),41,42,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 41 and dependent claims thereof

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

19. Claims: 1-6(in part),43,44,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 43 and dependent claims thereof

20. Claims: 1-6(in part),45,46,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 45 and dependent claims thereof

21. Claims: 1-6(in part),47,48,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 47 and dependent claims thereof

22. Claims: 1-6(in part),49,50,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 49 and dependent claims thereof

23. Claims: 1-6(in part),51,52,54(in part),56-57(in part)

Compounds further designated in claim 1, corresponding to compounds claimed in claim 51 and dependent claims thereof

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.1

Although claim 55 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box I.1

Claims Nos.: 53

The compounds of claim 53 lack to be properly defined: firstly because they are defined by a result to be achieved (i.e. as trypsin-like serine protease enzyme inhibitors), secondly because the term "P1" is not specified in the claim and has no support in the description (i.e. the expression "inhibitor comprising a P1 group" is not searchable as such).

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/20962

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9828326	A	02-07-1998	AU 5526098 A WO 9828326 A1	17-07-1998 02-07-1998
WO 9748706	A	24-12-1997	AU 3232597 A WO 9748706 A1 US 6124291 A	07-01-1998 24-12-1997 26-09-2000
WO 9619483	A	27-06-1996	AU 699679 B2 AU 4062795 A AU 715378 B2 AU 4062895 A AU 4250596 A AU 4250896 A BG 101647 A BR 9510433 A CA 2208772 A1 CA 2208773 A1 WO 9619483 A1 WO 9619491 A1 CN 1175259 A CZ 9701899 A3 EE 9700113 A EP 0802916 A1 EP 0799240 A1 FI 972466 A HU 77651 A2 JP 11508535 T JP 10513151 T LT 97132 A ,B LV 12019 A LV 12019 B MD 970253 A NO 972892 A NZ 297360 A PL 320965 A1 SK 83897 A3 ZA 9510960 A ZA 9510961 A	10-12-1998 27-06-1996 03-02-2000 04-07-1996 10-07-1996 10-07-1996 31-03-1998 10-11-1998 27-06-1996 27-06-1996 27-06-1996 27-06-1996 04-03-1998 16-09-1998 15-12-1997 29-10-1997 08-10-1997 19-08-1997 28-07-1998 27-07-1999 15-12-1998 25-03-1998 20-04-1998 20-07-1998 31-05-1999 20-08-1997 27-03-2000 24-11-1997 06-05-1998 09-07-1996 09-07-1996
WO 9857937	A	23-12-1998	AU 8150398 A BR 9810151 A EE 9900584 A EP 0991625 A2 HR 980334 A1 HU 0003906 A2 JP 2002507968 T LT 99146 A ,B LV 12516 A LV 12516 B NO 996316 A PL 337831 A1 SI 20208 A SK 174699 A3 WO 9857937 A2 ZA 9805251 A US 5998424 A	04-01-1999 08-08-2000 15-08-2000 12-04-2000 30-04-1999 28-05-2001 12-03-2002 25-05-2000 20-07-2000 20-03-2001 17-12-1999 11-09-2000 31-10-2000 14-08-2000 23-12-1998 17-12-1999 07-12-1999
WO 9723212	A	03-07-1997	AU 1335897 A	17-07-1997

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/20962

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9723212	A	CA 2240946 A1	03-07-1997
		EP 0874629 A1	04-11-1998
		HR 960597 A1	30-04-1998
		JP 2001502655 T	27-02-2001
		WO 9723212 A1	03-07-1997
		US 5939418 A	17-08-1999
		ZA 9610704 A	19-06-1998
